

1

SAMHSA Definition

Individual trauma results from an event, series of events, or set circumstances that is experienced by an individual as physically or emotionally harmful or life threatening and that has lasting adverse effects on the individual's functioning and mental, physical, social, emotional, or spiritual well-being.





etter Care. Better Patient Outcomes. Better Resul

2

Definitions

- · Secondary trauma
 - Results from exposure to another individual's traumatic experiences, rather than from exposure directly to a traumatic event
- Trauma-informed
 - Understanding trauma and awareness of its impact across settings, services, and populations
- Trauma-informed care
 - Strengths-based service delivery approach grounded in an understanding of and responsiveness to the impact of trauma



Better Care. Better Patient Outcomes. Better Resu

Keep in Mind...

- TIC is not necessarily therapeutic in nature
- Its main function is not intended to alleviate symptoms of toxic stress, but instead is intended to inform practice, procedure, and policy



Select

etter Care. Better Patient Outcomes. Better Result

4

Paradigm Shift

Trauma-informed care seeks to change the illness paradigm from one that asks, "What's wrong with you?" to, "What has happened to you?" (SAMHSA, 2015)

"A non-trauma-informed system punishes and blames your adult actions and asks, 'what's wrong with you?' A trauma-informed provider will hold you accountable for your adult actions but give you space and time to process 'what happened to you?' without adding guilt and more trauma."



Better Care. Better Patient Outcomes. Better Resul

5



(SAMHSA, 2015) Key Principles 1) Safety 2) Trustworthiness and Transparency 3) Peer Support 4) Collaboration Mutuality 5) Empowerment, Voice, Choice 6) Cultural, Historical, Gender Issues (SAMHSA, 2015)

7



8

Physical Safety Ensuring individuals feel they're being protected and that their physical safety is a priority Communicating clearly to staff, patients and community partners about crisis response efforts Soliciting input and feedback from staff Attending to staff unease

Select Retroduktorion Better Care. Better Patient Outcomes. Better Resul

Emotional Safety

- Helping staff understand what to expect
- Demonstrating flexible consistency
- Paying attention to nonverbal communication
- Building in time to check in about feelings



Select

ottor Caro Rottor Patient Outcomer, Rottor Recult

10

Trustworthiness and Transparency Organizational operations and decisions are conducted with transparency and the goal of building and maintaining trust among staff, clients, and family members of those receiving services

11

Select



Peer Support The term "Peer" refers to individuals with lived experiences of trauma Mutual self-help Vehicle for building trust, establishing safety, and empowerment

13

Peer Support and Relationship Supporting multiple ways for communicating Encouraging opportunities and methods for virtual face to face contact Setting up and supporting regular peer check-ins to connect Peer Support and Relationship Create a culture of calm "Physical distancing" not "social distancing"

14

Collaboration and Mutuality • Leveling power differences • Everyone is equal and has a role on the team • Healing happens in relationships and in the meaningful sharing of power and decision-making "One does not have to be a therapist to be therapeutic."

Collaboration

- Making your policy decisions with other systems in mind
- Working together to create policies that promote shared delivery of services, and working to eliminate policies where shared delivery is hindered
- Initiating new partnerships and nontraditional collaborations.
- Collaborating within the organization





Rotter Care Rotter Patient Outcomer, Rotter Recult

16

Empowerment, Voice and Choice

- Individual strengths are recognized
- Clients are supported in shared decision-making, choice, and goal setting to determine the plan of action.
- Cultivate self-advocacy skills
- Staff are facilitators of recovery rather than controllers of recovery





ter Care. Better Patient Outcomes. Better Result

17

Voice, Choice, and Empowerment



- Sharing power
- Providing choice whenever possible
- Providing staff with the scripts needed to explain the situation and policies to service users
- Listening to staff ideas and input about being trauma informed during this crisis

Select

etter Care. Better Patient Outcomes. Better Resul

Cultural, Historical, and Gender Issues

- Actively move past cultural stereotypes and biases
- Leverages the healing value of traditional cultural connections
- Incorporates policies, protocols, and processes that are responsive to the racial, ethnic and cultural needs of individuals served



Better Care, Better Patient Outcomes, Better Results

19

Cultural Responsivity

- Recognizing and building upon cultural strengths
- Ensuring intended and unintended consequences of policy and practice decisions do not create harm
- Being mindful of historical contexts
- Using strategies that encourage engagement and minimize mictures.
- Ensure important information is offered in multiple languages



etter Care. Better Patient Outcomes. Better Result

20

Context of Community

- Trauma does not occur in a vacuum
- Community can be defined:
 - · Geographically
 - Virtually
 - Organizationally

How a community responds to individual trauma sets the foundation for the impact of the traumatic event, experience and effect



Better Care. Better Patient Outcomes. Better Resu

Community Response to Trauma

- Communities that provide a context of understanding and selfdetermination
- Facilitate healing and recovery
- Communities that avoid, overlook or misunderstand the impact of trauma
 - May re-traumatize and interfere with the healing process



letter Care. Better Patient Outcomes. Better Results

22

Consider the Following...

- A community may be subjected to a community-threatening event
- Trauma can be transmitted from one generation to the next
- When explaining trauma-informed approaches, use language that reduces stigma
- Accommodate low health literacy
- Focus on how trauma affects health, not just the traumatic event/experience



etter Care. Better Patient Outcomes. Better Result

23

Impact of Trauma on Work

- Increased absenteeism from health problems or lost productivity
- Impaired judgement
- · Unwillingness to accept extra work or responsibility
- Low motivation
- Low productivity; poor work quality (leading to poor client outcomes)
- Decreased compliance with organizational requirements
- Staff conflict
- High staff turnover (leading to ongoing training costs)




Better Care. Better Patient Outcomes. Better Resu

Secondary Traumatization Signs Psychological distress Distressing emotions Nightmares, flooding, flashbacks of client disclosures Numbing or avoidance Somatic issues Addictive/compulsive behaviors Impaired functioning

25

Secondary Traumatization Signs Cognitive shifts Chronic suspicion about others Heightened sense of vulnerability Extreme sense of helplessness or exaggerated sense of control over others or situations Loss of personal control or freedom Bitterness or cynicism Blaming the victim or seeing everyone as a victim Witness or clinician guilt if client reexperiences trauma or reenacts trauma in counseling Feeling victimized by client



Secondary Traumatization Signs

Frame of reference

- Disconnection from one's sense of identity
- Dramatic change in fundamental beliefs about the world
- Loss or distortion of values or principles
- A previous sense of spirituality as comfort or resource decreases or becomes nonexistent
- · Loss of faith in something greater
- Existential despair and loneliness



28

Using Psychometric Measures

- If you use the ProQOL in clinical supervision, present it as a selfassessment tool
- Work collaboratively and respectfully to explore their own understanding of and meanings attached to their scores
- Scores are one way for you and your supervisees to get a sense of whether they might be at risk for secondary traumatization, what they can do to prevent it, how to address it, and how you can support them

ProQOL: A validated tool



29

Addressing Secondary Trauma

- · Engage in regular screening/self-
- assessment

 Address signs of STS within clinical supervision

 • Work collaboratively to develop a
- comprehensive self-care plan and evaluate its effectiveness on a regular basis
- Provide a safe and nonjudgmental environment
- Provide a place to debrief critical stress incidents at work Support and encourage individual
- counseling or psychotherapy, when needed

| Select | |
|----------------|--|
| Rehabilitation | |

Trauma Informed Leadership

- A way of understating or appreciating there is an emotional world of experiences in our colleagues
- Recognizes and honors the emotional scars that people may struggle with
- Helps leaders have empathy and compassion for their employees

Select Rehobilitation

Name of the Barbara Barbara Contraction Barbara Barbara

31

Why Trauma Informed Supervision

- Promotes staff retention and reduces turnover (Barak, et al., 2001; Knudsen et al., 2013)
- Reduces levels of vicarious trauma experienced by staff (WCSAP, 2004)
- Influences supervisee ability to more effectively cope and have resilience (Sommer & Cox, 2005; Turner, 2009)
- Enhances worker well-being
- Facilitates best practices (Schwalbe et al., 2014)



etter Care. Better Patient Outcomes. Better Result

32

What is Trauma Informed Supervision?

- Supporting staff
- Relationship based supervision
- Best practices for reflective supervision
 - Regularly scheduled meetings without interruption
 - Discussions about change management
 - Inviting colleagues to share experiences and explore experiences



Better Care. Better Patient Outcomes. Better Resu

Leading with Empathy

When employees experience empathy, understanding and compassion from the leadership team they will feel safe, respected and recognized



etter Care. Better Patient Outcomes. Better Results.

34


Staff Responses

- Stress will transfer to our residents
- Give yourself some space to cope
- Share resources and education about abuse
- Be watchful and vigilant for risk factors for suicide
- Practice self-kindness
- Take care of yourself
- Gain insight into where you are struggling



tter Care. Better Patient Outcomes. Better Result

35





37

Negative and Unrealistic Thinking • All-or-nothing thinking • Overgeneralization • The mental filter • Diminishing the positive • Jumping to conclusions • Emotional reasoning • 'Shoulds' and 'should-nots' • Labeling

38

The Witness Stand "What's the evidence that this thought is true? Not true?" "What would I tell a friend who had this thought?" "Is there another way of looking at the situation or an alternate explanation?" "How might I look at this situation if I didn't have depression?"

During work shifts: Self-monitoring and pacing Check-ins with colleagues, family, and friends Working in partnerships or in teams Brief relaxation/stress management breaks Time-outs Keep anxieties conscribed to actual threats Helpful self-talk Focus efforts on what is within your power Accept what you cannot change

40

Strategies to Help Residents Establishing a routine and maintaining clear communication are crucial Do not assume residents understand why the routine is changing Do not assume residents' understanding of a global pandemic are universal Relationships and well-being can take priority Create relational rituals

41

Select

Supporting Safety Offer residents a way to connect if there is something that they need help with or are worried about Encourage residents to talk to friends or family members on the phone Help residents plan virtual get togethers with family Keep as much of a regular routine as possible Plan fun activities

Connectedness

- Make time to ask residents about something they are doing right now
- Plan activities through the use of web-conferencing sites
- Write letters
- Talk directly about the importance of connecting with others

"Physical distancing" NOT "social distancing"





Better Care. Better Patient Outcomes. Better Result

43

Hope

- Have residents connect with someone in their family or community to ask another person they respect how they stayed hopeful in troubled times
- Download and view YouTube videos of church or inspirational speakers
- Take time to learn about other historical times of crisis, including how these ended and communities rebounded
- Encourage residents to get fresh air and to move when possible
- Share some of the many stories of hope and helping that have come out of this current crisis

Hope is being able to see that there is light despite all of the darkness.

Desmond Tutu



Better Care. Better Patient Outcomes. Better Resul

44

What Else??

- Provide opportunities for residents to engage in activities that
 affirm their competence, sense of self-worth and feelings of
 safety
- Promote self-awareness
- Mindfulness or self-soothing exercises
- Model and normalize a range of emotions by giving residents opportunities to express themselves in nonverbal ways
- Virtual or long-distance appreciation or gratitude circle



Better Care. Better Patient Outcomes. Better Resul

At Risk Individuals

- Residents who have anxiety
- Residents who have depression or suicidal ideation
- Residents whose families may have lost jobs or income
- Residents who have loved ones particularly vulnerable to the COVID-19 virus
- Residents who have a caregiver who is a healthcare worker or in another occupation where they are exposed to the virus or are being asked to respond in an intense way



46

Considerations

- Support regulationPrioritize relationships
- Explain the why behind decisions
- Help staff know what to expect to the extent possible
- Reframe behaviors



47

Remember that we as caregivers are the best predictors of how our residents are doing; they are watching and listening to us. When we take care of ourselves, we're showing them how they can take care of themselves, too.

48

Select